

Mathematical Logic

Practical Class: Set Theory

Chiara Ghidini

FBK-IRST, Trento, Italy

2014/2015

1 Set Theory

- Basic Concepts
- Operations on Sets
- Operation Properties

2 Relations

- Properties
- Equivalence Relation

3 Functions


- Properties

Sets: Basic Concepts

- The concept of **set** is considered a primitive concept in math
- A set is a collection of elements whose description must be unambiguous and unique: it must be possible to decide whether an element belongs to the set or not.
- **Examples:**
 - *the students in this classroom*
 - *the points in a straight line*
 - *the cards in a playing pack*
- are all sets, while
 - *students that hates math*
 - *amusing books*
- are not sets.

Describing Sets

- In set theory there are several description methods:
 - **Listing**: the set is described listing all its elements
Example: $A = \{a, e, i, o, u\}$.
 - **Abstraction**: the set is described through a property of its elements
Example: $A = \{x \mid x \text{ is a vowel of the Latin alphabet}\}$.
 - **Eulero-Venn Diagrams**: graphical representation that supports the formal description



Sets: Basic Concepts (2)

- **Empty Set:** \emptyset , is the set containing no elements;
- **Membership:** $a \in A$, element a belongs to the set A ;
 - *Non membership:* $a \notin A$, element a doesn't belong to the set A ;
- **Equality:** $A = B$, iff the sets A and B contain the same elements;
 - *inequality:* $A \neq B$, iff it is not the case that $A = B$;
- **Subset:** $A \subseteq B$, iff all elements in A belong to B too;
- **Proper subset:** $A \subset B$, iff $A \subseteq B$ and $A \neq B$.

Power set

- We define the **power set** of a set A , denoted with $P(A)$, as the set containing all the subsets of A .

Power set

- We define the **power set** of a set A , denoted with $P(A)$, as the set containing all the subsets of A .
- **Example:** if $A = \{a, b, c\}$, then
$$P(A) = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}, \}$$

Power set

- We define the **power set** of a set A , denoted with $P(A)$, as the set containing all the subsets of A .
- **Example:** if $A = \{a, b, c\}$, then
$$P(A) = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}, \}$$
- If A has n elements, then its power set $P(A)$ contains 2^n elements.
 - Exercise: prove it!!!

Operations on Sets

- **Union**: given two sets A and B we define the union of A and B as the set containing the elements belonging to A or to B or to both of them, and we denote it with $A \cup B$.

Operations on Sets

- **Union:** given two sets A and B we define the union of A and B as the set containing the elements belonging to A or to B or to both of them, and we denote it with $A \cup B$.
 - **Example:** if $A = \{a, b, c\}$, $B = \{a, d, e\}$ then
 $A \cup B = \{a, b, c, d, e\}$

Operations on Sets

- **Union:** given two sets A and B we define the union of A and B as the set containing the elements belonging to A or to B or to both of them, and we denote it with $A \cup B$.
 - **Example:** if $A = \{a, b, c\}$, $B = \{a, d, e\}$ then
$$A \cup B = \{a, b, c, d, e\}$$
- **Intersection:** given two sets A and B we define the intersection of A and B as the set containing the elements that belongs both to A and B , and we denote it with $A \cap B$.

Operations on Sets

- **Union:** given two sets A and B we define the union of A and B as the set containing the elements belonging to A or to B or to both of them, and we denote it with $A \cup B$.
 - **Example:** if $A = \{a, b, c\}$, $B = \{a, d, e\}$ then
 $A \cup B = \{a, b, c, d, e\}$
- **Intersection:** given two sets A and B we define the intersection of A and B as the set containing the elements that belongs both to A and B , and we denote it with $A \cap B$.
 - **Example:** if $A = \{a, b, c\}$, $B = \{a, d, e\}$ then
 $A \cap B = \{a\}$

Operations on Sets (2)

- **Difference:** given two sets A and B we define the difference of A and B as the set containing all the elements which are members of A , but not members of B , and denote it with $A - B$.

Operations on Sets (2)

- **Difference:** given two sets A and B we define the difference of A and B as the set containing all the elements which are members of A , but not members of B , and denote it with $A - B$.
 - **Example:** if $A = \{a, b, c\}$, $B = \{a, d, e\}$ then $A - B = \{b, c\}$

Operations on Sets (2)

- **Difference:** given two sets A and B we define the difference of A and B as the set containing all the elements which are members of A , but not members of B , and denote it with $A - B$.
 - **Example:** if $A = \{a, b, c\}$, $B = \{a, d, e\}$ then $A - B = \{b, c\}$
- **Complement:** given a universal set U and a set A , where $A \subseteq U$, we define the complement of A in U , denoted with \overline{A} (or $C_U A$), as the set containing all the elements in U not belonging to A .

Operations on Sets (2)

- **Difference:** given two sets A and B we define the difference of A and B as the set containing all the elements which are members of A , but not members of B , and denote it with $A - B$.
 - **Example:** if $A = \{a, b, c\}$, $B = \{a, d, e\}$ then $A - B = \{b, c\}$
- **Complement:** given a universal set U and a set A , where $A \subseteq U$, we define the complement of A in U , denoted with \overline{A} (or $C_U A$), as the set containing all the elements in U not belonging to A .
 - **Example:** if U is the set of natural numbers and A is the set of even numbers (0 included), then the complement of A in U is the set of odd numbers.

Sets: Examples

- **Examples:**

- Given $A = \{a, e, i, o, \{u\}\}$ and $B = \{i, o, u\}$, consider the following statements:

- 1 $B \in A$

Sets: Examples

- **Examples:**

- Given $A = \{a, e, i, o, \{u\}\}$ and $B = \{i, o, u\}$, consider the following statements:

① $B \in A$ **NO!**

Sets: Examples

- **Examples:**

- Given $A = \{a, e, i, o, \{u\}\}$ and $B = \{i, o, u\}$, consider the following statements:

① $B \in A$ **NO!**

② $(B - \{i, o\}) \in A$

Sets: Examples

- **Examples:**

- Given $A = \{a, e, i, o, \{u\}\}$ and $B = \{i, o, u\}$, consider the following statements:

① $B \in A$ **NO!**

② $(B - \{i, o\}) \in A$ **OK**

Sets: Examples

- **Examples:**

- Given $A = \{a, e, i, o, \{u\}\}$ and $B = \{i, o, u\}$, consider the following statements:

① $B \in A$ **NO!**

② $(B - \{i, o\}) \in A$ **OK**

③ $\{a\} \cup \{i\} \subset A$

Sets: Examples

- **Examples:**

- Given $A = \{a, e, i, o, \{u\}\}$ and $B = \{i, o, u\}$, consider the following statements:

- 1 $B \in A$ **NO!**
- 2 $(B - \{i, o\}) \in A$ **OK**
- 3 $\{a\} \cup \{i\} \subset A$ **OK**

Sets: Examples

- **Examples:**

- Given $A = \{a, e, i, o, \{u\}\}$ and $B = \{i, o, u\}$, consider the following statements:

- 1 $B \in A$ **NO!**
- 2 $(B - \{i, o\}) \in A$ **OK**
- 3 $\{a\} \cup \{i\} \subset A$ **OK**
- 4 $\{u\} \subset A$

Sets: Examples

- **Examples:**

- Given $A = \{a, e, i, o, \{u\}\}$ and $B = \{i, o, u\}$, consider the following statements:

- ① $B \in A$ **NO!**
- ② $(B - \{i, o\}) \in A$ **OK**
- ③ $\{a\} \cup \{i\} \subset A$ **OK**
- ④ $\{u\} \subset A$ **NO!**

Sets: Examples

- **Examples:**

- Given $A = \{a, e, i, o, \{u\}\}$ and $B = \{i, o, u\}$, consider the following statements:

- 1 $B \in A$ **NO!**
- 2 $(B - \{i, o\}) \in A$ **OK**
- 3 $\{a\} \cup \{i\} \subset A$ **OK**
- 4 $\{u\} \subset A$ **NO!**
- 5 $\{\{u\}\} \subset A$

Sets: Examples

- **Examples:**

- Given $A = \{a, e, i, o, \{u\}\}$ and $B = \{i, o, u\}$, consider the following statements:

- 1 $B \in A$ **NO!**
- 2 $(B - \{i, o\}) \in A$ **OK**
- 3 $\{a\} \cup \{i\} \subset A$ **OK**
- 4 $\{u\} \subset A$ **NO!**
- 5 $\{\{u\}\} \subset A$ **OK**

Sets: Examples

- **Examples:**

- Given $A = \{a, e, i, o, \{u\}\}$ and $B = \{i, o, u\}$, consider the following statements:

- 1 $B \in A$ **NO!**
- 2 $(B - \{i, o\}) \in A$ **OK**
- 3 $\{a\} \cup \{i\} \subset A$ **OK**
- 4 $\{u\} \subset A$ **NO!**
- 5 $\{\{u\}\} \subset A$ **OK**
- 6 $B - A = \emptyset$

Sets: Examples

- **Examples:**

- Given $A = \{a, e, i, o, \{u\}\}$ and $B = \{i, o, u\}$, consider the following statements:

① $B \in A$ **NO!**

② $(B - \{i, o\}) \in A$ **OK**

③ $\{a\} \cup \{i\} \subset A$ **OK**

④ $\{u\} \subset A$ **NO!**

⑤ $\{\{u\}\} \subset A$ **OK**

⑥ $B - A = \emptyset$ **NO!** $B - A = \{u\}$

Sets: Examples

- **Examples:**

- Given $A = \{a, e, i, o, \{u\}\}$ and $B = \{i, o, u\}$, consider the following statements:

① $B \in A$ **NO!**

② $(B - \{i, o\}) \in A$ **OK**

③ $\{a\} \cup \{i\} \subset A$ **OK**

④ $\{u\} \subset A$ **NO!**

⑤ $\{\{u\}\} \subset A$ **OK**

⑥ $B - A = \emptyset$ **NO!** $B - A = \{u\}$

⑦ $i \in A \cap B$

Sets: Examples

- **Examples:**

- Given $A = \{a, e, i, o, \{u\}\}$ and $B = \{i, o, u\}$, consider the following statements:

① $B \in A$ **NO!**

② $(B - \{i, o\}) \in A$ **OK**

③ $\{a\} \cup \{i\} \subset A$ **OK**

④ $\{u\} \subset A$ **NO!**

⑤ $\{\{u\}\} \subset A$ **OK**

⑥ $B - A = \emptyset$ **NO!** $B - A = \{u\}$

⑦ $i \in A \cap B$ **OK**

Sets: Examples

- **Examples:**

- Given $A = \{a, e, i, o, \{u\}\}$ and $B = \{i, o, u\}$, consider the following statements:

① $B \in A$ **NO!**

② $(B - \{i, o\}) \in A$ **OK**

③ $\{a\} \cup \{i\} \subset A$ **OK**

④ $\{u\} \subset A$ **NO!**

⑤ $\{\{u\}\} \subset A$ **OK**

⑥ $B - A = \emptyset$ **NO!** $B - A = \{u\}$

⑦ $i \in A \cap B$ **OK**

⑧ $\{i, o\} = A \cap B$

Sets: Examples

- **Examples:**

- Given $A = \{a, e, i, o, \{u\}\}$ and $B = \{i, o, u\}$, consider the following statements:

① $B \in A$ **NO!**

② $(B - \{i, o\}) \in A$ **OK**

③ $\{a\} \cup \{i\} \subset A$ **OK**

④ $\{u\} \subset A$ **NO!**

⑤ $\{\{u\}\} \subset A$ **OK**

⑥ $B - A = \emptyset$ **NO!** $B - A = \{u\}$

⑦ $i \in A \cap B$ **OK**

⑧ $\{i, o\} = A \cap B$ **OK**

Sets: Exercises

- Exercises:

- Given $A = \{t, z\}$ and $B = \{v, z, t\}$ consider the following statements:

- 1 $A \in B$

- 2 $A \subset B$

- 3 $z \in A \cap B$

- 4 $v \subset B$

- 5 $\{v\} \subset B$

- 6 $v \in A - B$

- Given $A = \{a, b, c, d\}$ and $B = \{c, d, f\}$

- find a set X s.t. $A \cup B = B \cup X$; is this set unique?
- there exists a set Y s.t. $A \cup Y = B$?

Sets: Exercises (2)

- Exercises:

- Given $A = \{0, 2, 4, 6, 8, 10\}$, $B = \{0, 1, 2, 3, 4, 5, 6\}$ and $C = \{4, 5, 6, 7, 8, 9, 10\}$, compute:
 - $A \cap B \cap C$, $A \cup (B \cap C)$, $A - (B - C)$
 - $(A \cup B) \cap C$, $(A - B) - C$, $A \cap (B - C)$
- Describe 3 sets A, B, C s.t. $A \cap (B \cup C) \neq (A \cap B) \cup C$

Sets: Operation Properties

- $A \cap A = A,$
 $A \cup A = A$

Sets: Operation Properties

- $A \cap A = A,$
 $A \cup A = A$
- $A \cap B = B \cap A,$
 $A \cup B = B \cup A$ (*commutative*)

Sets: Operation Properties

- $A \cap A = A,$
 $A \cup A = A$
- $A \cap B = B \cap A,$
 $A \cup B = B \cup A$ (*commutative*)
- $A \cap \emptyset = \emptyset,$
 $A \cup \emptyset = A$

Sets: Operation Properties

- $A \cap A = A,$
 $A \cup A = A$
- $A \cap B = B \cap A,$
 $A \cup B = B \cup A$ (*commutative*)
- $A \cap \emptyset = \emptyset,$
 $A \cup \emptyset = A$
- $(A \cap B) \cap C = A \cap (B \cap C),$
 $(A \cup B) \cup C = A \cup (B \cup C)$ (*associative*)

Sets: Operation Properties(2)

- $A \cap (B \cup C) = (A \cap B) \cup (A \cap C),$
 $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
(*distributive*)

Sets: Operation Properties(2)

- $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$,
 $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
(*distributive*)
- $\overline{A \cap B} = \overline{A} \cup \overline{B}$,
 $\overline{A \cup B} = \overline{A} \cap \overline{B}$ (*De Morgan laws*)

Sets: Operation Properties(2)

- $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$,
 $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
(*distributive*)
- $\overline{A \cap B} = \overline{A} \cup \overline{B}$,
 $\overline{A \cup B} = \overline{A} \cap \overline{B}$ (*De Morgan laws*)
- **Exercise:** Prove the validity of all the properties.

Cartesian Product

- Given two sets A and B , we define the **Cartesian product** of A and B as the set of ordered couples (a, b) where $a \in A$ and $b \in B$; formally,

$$A \times B = \{(a, b) : a \in A \text{ and } b \in B\}$$

- Notice that: $A \times B \neq B \times A$

Cartesian Product (2)

- **Examples:**

- given $A = \{1, 2, 3\}$ and $B = \{a, b\}$, then

$$A \times B = \{(1, a), (1, b), (2, a), (2, b), (3, a), (3, b)\} \text{ and}$$

$$B \times A = \{(a, 1), (a, 2), (a, 3), (b, 1), (b, 2), (b, 3)\}.$$

Cartesian Product (2)

- **Examples:**

- given $A = \{1, 2, 3\}$ and $B = \{a, b\}$, then
 $A \times B = \{(1, a), (1, b), (2, a), (2, b), (3, a), (3, b)\}$ and
 $B \times A = \{(a, 1), (a, 2), (a, 3), (b, 1), (b, 2), (b, 3)\}$.
- Cartesian coordinates of the points in a plane are an example of the Cartesian product $\mathfrak{R} \times \mathfrak{R}$

Cartesian Product (2)

- **Examples:**

- given $A = \{1, 2, 3\}$ and $B = \{a, b\}$, then
 $A \times B = \{(1, a), (1, b), (2, a), (2, b), (3, a), (3, b)\}$ and
 $B \times A = \{(a, 1), (a, 2), (a, 3), (b, 1), (b, 2), (b, 3)\}$.
- Cartesian coordinates of the points in a plane are an example of the Cartesian product $\mathfrak{R} \times \mathfrak{R}$
- The Cartesian product can be computed on any number n of sets A_1, A_2, \dots, A_n , $A_1 \times A_2 \times \dots \times A_n$ is the set of ordered n -tuple (x_1, \dots, x_n) where $x_i \in A_i$ for each $i = 1 \dots n$.

Relations

- A **relation** R from the set A to the set B is a subset of the Cartesian product of A and B : $R \subseteq A \times B$; if $(x, y) \in R$, then we will write xRy for 'x is R -related to y '.

Relations

- A **relation** R from the set A to the set B is a subset of the Cartesian product of A and B : $R \subseteq A \times B$; if $(x, y) \in R$, then we will write xRy for 'x is R -related to y '.
- A binary relation on a set A is a subset $R \subseteq A \times A$

Relations

- A **relation** R from the set A to the set B is a subset of the Cartesian product of A and B : $R \subseteq A \times B$; if $(x, y) \in R$, then we will write xRy for ' x is R -related to y '.
- A binary relation on a set A is a subset $R \subseteq A \times A$
- **Examples:**
 - given $A = \{1, 2, 3, 4\}$, $B = \{a, b, d, e, r, t\}$ and aRb iff in the Italian name of a there is the letter b , then
$$R = \{(2, d), (2, e), (3, e), (3, r), (3, t), (4, a), (4, r), (4, t)\}$$

Relations

- A **relation** R from the set A to the set B is a subset of the Cartesian product of A and B : $R \subseteq A \times B$; if $(x, y) \in R$, then we will write xRy for 'x is R -related to y '.
- A binary relation on a set A is a subset $R \subseteq A \times A$
- **Examples:**
 - given $A = \{1, 2, 3, 4\}$, $B = \{a, b, d, e, r, t\}$ and aRb iff in the Italian name of a there is the letter b , then
$$R = \{(2, d), (2, e), (3, e), (3, r), (3, t), (4, a), (4, r), (4, t)\}$$
 - given $A = \{3, 5, 7\}$, $B = \{2, 4, 6, 8, 10, 12\}$ and aRb iff a is a divisor of b , then
$$R = \{(3, 6), (3, 12), (5, 10)\}$$

Relations

- A **relation** R from the set A to the set B is a subset of the Cartesian product of A and B : $R \subseteq A \times B$; if $(x, y) \in R$, then we will write xRy for 'x is R -related to y '.
- A binary relation on a set A is a subset $R \subseteq A \times A$
- **Examples:**
 - given $A = \{1, 2, 3, 4\}$, $B = \{a, b, d, e, r, t\}$ and aRb iff in the Italian name of a there is the letter b , then
$$R = \{(2, d), (2, e), (3, e), (3, r), (3, t), (4, a), (4, r), (4, t)\}$$
 - given $A = \{3, 5, 7\}$, $B = \{2, 4, 6, 8, 10, 12\}$ and aRb iff a is a divisor of b , then
$$R = \{(3, 6), (3, 12), (5, 10)\}$$
- **Exercise:** in prev example, let aRb iff $a + b$ is an even number
 $R = ?$

Relations (2)

- Given a relation R from A to B ,
 - the **domain** of R is the set $Dom(R) = \{a \in A \mid \text{there exists a } b \in B, aRb\}$
 - the **co-domain** of R is the set $Cod(R) = \{b \in B \mid \text{there exists an } a \in A, aRb\}$

Relations (2)

- Given a relation R from A to B ,
 - the **domain** of R is the set $Dom(R) = \{a \in A \mid \text{there exists a } b \in B, aRb\}$
 - the **co-domain** of R is the set $Cod(R) = \{b \in B \mid \text{there exists an } a \in A, aRb\}$
- Let R be a relation from A to B . The **inverse relation** of R is the relation $R^{-1} \subseteq B \times A$ where $R^{-1} = \{(b, a) \mid (a, b) \in R\}$

Relation properties

- Let R be a binary relation on A . R is
 - **reflexive** iff aRa for all $a \in A$;
 - **symmetric** iff aRb implies bRa for all $a, b \in A$;
 - **transitive** iff aRb and bRc imply aRc for all $a, b, c \in A$;
 - **anti-symmetric** iff aRb and bRa imply $a = b$ for all $a, b \in A$;

Equivalence Relation

- Let R be a binary relation on a set A . R is an **equivalence relation** iff it satisfies all the following properties:
 - reflexive
 - symmetric
 - transitive
- an equivalence relation is usually denoted with \sim or \equiv

Set Partition

- Let A be a set, a **partition** of A is a family F of non-empty subsets of A s.t.:
 - the subsets are pairwise disjoint
 - the union of all the subsets is the set A
- Notice that: each element of A belongs to exactly one subset in F .

Equivalence Classes

- Let A be a set and \equiv an equivalence relation on A , given an $x \in A$ we define **equivalence class** X the set of elements $x' \in A$ s.t. $x' \equiv x$, formally
$$X = \{x' \mid x' \equiv x\}$$

Equivalence Classes

- Let A be a set and \equiv an equivalence relation on A , given an $x \in A$ we define **equivalence class** X the set of elements $x' \in A$ s.t. $x' \equiv x$, formally
$$X = \{x' \mid x' \equiv x\}$$
- Notice that: any element x is sufficient to obtain the equivalence class X , which is denoted also with $[x]$
 - $x \equiv x'$ implies $[x] = [x'] = X$

Equivalence Classes

- Let A be a set and \equiv an equivalence relation on A , given an $x \in A$ we define **equivalence class** X the set of elements $x' \in A$ s.t. $x' \equiv x$, formally
$$X = \{x' \mid x' \equiv x\}$$
- Notice that: any element x is sufficient to obtain the equivalence class X , which is denoted also with $[x]$
 - $x \equiv x'$ implies $[x] = [x'] = X$
- We define **quotient set** of A with respect to an equivalence relation \equiv as the set of equivalence classes defined by \equiv on A , and denote it with A/\equiv

Equivalence Classes (2)

- **Theorem:** Given an equivalence relation \equiv on A , the equivalence classes defined by \equiv on A are a partition of A . Similarly, given a partition on A , the relation R defined as xRx' iff x and x' belong to the same subset, is an equivalence relation on A .

Equivalence classes (3)

- **Example:** Parallelism relation.
Two straight lines in a plane are parallel if they do not have any point in common or if they coincide.

Equivalence classes (3)

- **Example:** Parallelism relation.
Two straight lines in a plane are parallel if they do not have any point in common or if they coincide.
- The parallelism relation \parallel is an equivalence relation since it is:

Equivalence classes (3)

- **Example:** Parallelism relation.
Two straight lines in a plane are parallel if they do not have any point in common or if they coincide.
- The parallelism relation \parallel is an equivalence relation since it is:
 - reflexive $r \parallel r$

Equivalence classes (3)

- **Example:** Parallelism relation.
Two straight lines in a plane are parallel if they do not have any point in common or if they coincide.
- The parallelism relation \parallel is an equivalence relation since it is:
 - reflexive $r \parallel r$
 - symmetric $r \parallel s$ implies $s \parallel r$

Equivalence classes (3)

- **Example:** Parallelism relation.
Two straight lines in a plane are parallel if they do not have any point in common or if they coincide.
- The parallelism relation \parallel is an equivalence relation since it is:
 - reflexive $r \parallel r$
 - symmetric $r \parallel s$ implies $s \parallel r$
 - transitive $r \parallel s$ and $s \parallel t$ imply $r \parallel t$

Equivalence classes (3)

- **Example:** Parallelism relation.
Two straight lines in a plane are parallel if they do not have any point in common or if they coincide.
- The parallelism relation \parallel is an equivalence relation since it is:
 - reflexive $r \parallel r$
 - symmetric $r \parallel s$ implies $s \parallel r$
 - transitive $r \parallel s$ and $s \parallel t$ imply $r \parallel t$
- We can thus obtain a partition in equivalence classes:
intuitively, each class represent a direction in the plane.

Order Relation

- Let A be a set and R be a binary relation on A . R is an **order** (partial), usually denoted with \leq , if it satisfies the following properties:
 - reflexive $a \leq a$
 - anti-symmetric $a \leq b$ and $b \leq a$ imply $a = b$
 - transitive $a \leq b$ and $b \leq c$ imply $a \leq c$

Order Relation

- Let A be a set and R be a binary relation on A . R is an **order** (partial), usually denoted with \leq , if it satisfies the following properties:
 - reflexive $a \leq a$
 - anti-symmetric $a \leq b$ and $b \leq a$ imply $a = b$
 - transitive $a \leq b$ and $b \leq c$ imply $a \leq c$
- If the relation holds for all $a, b \in A$ then it is a **total order**

Order Relation

- Let A be a set and R be a binary relation on A . R is an **order** (partial), usually denoted with \leq , if it satisfies the following properties:
 - reflexive $a \leq a$
 - anti-symmetric $a \leq b$ and $b \leq a$ imply $a = b$
 - transitive $a \leq b$ and $b \leq c$ imply $a \leq c$
- If the relation holds for all $a, b \in A$ then it is a **total order**
- A relation is a **strict order**, denoted with $<$, if it satisfies the following properties:
 - transitive $a < b$ and $b < c$ imply $a < c$
 - for all $a, b \in A$ either $a < b$ or $b < a$ or $a = b$

Relations : Exercises

- Exercises:
 - Decide whether the following relations $R : \mathbb{Z} \times \mathbb{Z}$ are symmetric, reflexive and transitive:
 - $R = \{(n, m) \in \mathbb{Z} \times \mathbb{Z} : n = m\}$
 - $R = \{(n, m) \in \mathbb{Z} \times \mathbb{Z} : |n - m| = 5\}$
 - $R = \{(n, m) \in \mathbb{Z} \times \mathbb{Z} : n \geq m\}$

Relations : Exercises (2)

- Exercises:

- Let $X = \{1, 2, 3, \dots, 30, 31\}$. Consider the relation on X : xRy if the dates x and y of January 2006 are on the same day of the week (Monday, Tuesday ..). Is R an equivalence relation? If this is the case describe its equivalence classes.
- Let $X = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$
 - Consider the following relation on X : xRy iff $x + y$ is an even number. Is R an equivalence relation? If this is the case describe its equivalence classes.
 - Consider the following relation on X : xRy iff $x + y$ is an odd number. Is R an equivalence relation? If this is the case describe its equivalence classes.

Relations : Exercises (3)

- Exercises:

- Let X be the set of straight-lines in the plane, and let x be a point in the plane. Are the following relations equivalence relations? If this is the case describe the equivalence classes.
 - $r \sim s$ iff r and s are parallel
 - $r \sim s$ iff the distance between r and x is equal to the distance between s and x
 - $r \sim s$ iff r and s are perpendicular
 - $r \sim s$ iff the distance between r and x is greater or equal to the distance between s and x
 - $r \sim s$ iff both r and s pass through x

Relations : Exercises (4)

- Exercises:

- Let div be a relation on \mathbb{N} defined as $a div b$ iff a divides b .
Where a divides b iff there exists an $n \in \mathbb{N}$ s.t. $a * n = b$
 - Is div an equivalence relation?
 - Is div an order?

Functions

- Given two sets A and B , a **function** f from A to B is a relation that associates to each element a in A exactly one element b in B . Denoted with $f : A \rightarrow B$

Functions

- Given two sets A and B , a **function** f from A to B is a relation that associates to each element a in A exactly one element b in B . Denoted with
 $f : A \longrightarrow B$
- The domain of f is the whole set A ; the image of each element a in A is the element b in B s.t. $b = f(a)$; the co-domain of f (or image of f) is a subset of B defined as follows:
 $Im_f = \{b \in B \mid \text{there exists an } a \in A \text{ s.t. } b = f(a)\}$

Functions

- Given two sets A and B , a **function** f from A to B is a relation that associates to each element a in A exactly one element b in B . Denoted with
$$f : A \longrightarrow B$$
- The domain of f is the whole set A ; the image of each element a in A is the element b in B s.t. $b = f(a)$; the co-domain of f (or image of f) is a subset of B defined as follows:
$$Im_f = \{b \in B \mid \text{there exists an } a \in A \text{ s.t. } b = f(a)\}$$
- Notice that: it can be the case that the same element in B is the image of several elements in A .

Classes of functions

- A function $f : A \rightarrow B$ is **surjective** if each element in B is image of some elements in A :
for each $b \in B$ there exists an $a \in A$ s.t. $f(a) = b$

Classes of functions

- A function $f : A \longrightarrow B$ is **surjective** if each element in B is image of some elements in A :
for each $b \in B$ there exists an $a \in A$ s.t. $f(a) = b$
- A function $f : A \longrightarrow B$ is **injective** if distinct elements in A have distinct images in B :
for each $b \in Im_f$ there exists a unique $a \in A$ s.t. $f(a) = b$

Classes of functions


- A function $f : A \rightarrow B$ is **surjective** if each element in B is image of some elements in A :
for each $b \in B$ there exists an $a \in A$ s.t. $f(a) = b$
- A function $f : A \rightarrow B$ is **injective** if distinct elements in A have distinct images in B :
for each $b \in \text{Im}_f$ there exists a unique $a \in A$ s.t. $f(a) = b$
- A function $f : A \rightarrow B$ is **bijective** if it is injective and surjective:
for each $b \in B$ there exists a unique $a \in A$ s.t. $f(a) = b$

Inverse Function

- If $f : A \longrightarrow B$ is bijective we can define its **inverse function**:
 $f^{-1} : B \longrightarrow A$

Inverse Function

- If $f : A \rightarrow B$ is bijective we can define its **inverse function**:
 $f^{-1} : B \rightarrow A$
- For each function f we can define its inverse relation; such a relation is a function iff f is bijective.
- Example:



the inverse relation of f is NOT a function.

Composed functions

- Let $f : A \rightarrow B$ and $g : B \rightarrow C$ be functions. The **composition** of f and g is the function $g \circ f : A \rightarrow C$ obtained by applying f and then g :
 $(g \circ f)(a) = g(f(a))$ for each $a \in A$
 $g \circ f = \{(a, g(f(a))) \mid a \in A\}$

Functions : Exercises

• Exercises:

- Given $A = \{ \text{students that passed the Logic exam} \}$ and $B = \{18, 19, \dots, 29, 30, 30L\}$, and let $f : A \rightarrow B$ be the function defined as $f(x) = \text{grade of } x \text{ in Logic}$. Answer the following questions:
 - What is the image of f ?
 - Is f bijective?
- Let A be the set of all people, and let $f : A \rightarrow A$ be the function defined as $f(x) = \text{father of } x$. Answer the following questions:
 - What is the image of f ?
 - Is f bijective?
 - Is f invertible?
- Let $f : \mathbb{N} \rightarrow \mathbb{N}$ be the function defined as $f(n) = 2n$.
 - What is the image of f ?
 - Is f bijective?
 - Is f invertible?